REFERENCE

Abida, Narasimhan.B, Srinivas.K, Imran.M,(2011), RP-HPLC method development and validation for gatifloxacin from tablet formulation, IJPI's J Anal Chem,2(7),1-8.

Aboul-Enein.H.Y, Hoenen, Ghanem.A, Koll.M,(2005), Reversed phase liquid chromatographic method for the high-throughput analysis of clopidogrel in pharmaceutical formulations using a monolithic silica column, J Liq.Chromato. RT,28(9),1357-1365.

Al-Dgither.S, Naseeruddin.A.S, Hammami,(2006), Development and validation of an HPLC method for the determination of gatifloxacin stability in human plasma, J Pharm.Biomed.Anal.,41,251-255.

Ashwini.E.P, Shila.V.D, Nayana.D.P, Sheetal.V.P, Manoj.M.B, Shashikant.D.B, (2013),Validated RP-HPLC method for simultaneous estimation of amlodipine besylate and clopidogrelbisulphate in bulk and tablet dosage form, Int. J Bioassays,2(2), 412-415.

Bhumika.D.S, Priti.D.T, Darshana.K.M,(2012), Development and validation of spectrophotometric methods for simultaneous estimation of escitalopram oxalate and etizolam in their combined tablet dosage form., J.Pharma.Sci.Biosci.Res., 2(5),195-200.

Borner.K, Hartwig.H, Lode.H,(2000), Determination of gatifloxacin in human serum and urine by HPLC, Chromatographia,52,S105-S107.

Carazos.M.L.S, Gonzalez.L.Y.C, Lerma.G.G, Torres.N.W,(2006), Determination of gatifloxacin in semen by HPLC with diode-array and fluorescence detection, Chromatographia,63,605-608.

Crevar-Sakac.M, Vujic.Z, Brboric.J, Kuntic.V, Uskokovic-Markovic,(2013), An improved HPLC method with the aid of a chemometric protocol: simultaneous determination of atorvastatin and its metabolites in plasma, Molecules, 18(3), 2469-2482.

Dillon.T, Bondarenko.P, Pipes.G, Ricci.M, Rehder.D, Kleemann.G,(2008), LC/MS method of analyzing high molecular weight proteins.US7329353 B2.

Erturk.S, Sevinc-Aktas.E, Ersoy.L, Ficicioglu.S,(2003), An HPLC method for the Biomed.Anal.,33,1017-1023.

Gandhi.D.S, Sabnis.B, Bothara.K,(2008), Simultaneous HPTLC determination of escitalopram oxalate and clonazepam in combined tablets, Chromatographia,67(5-6),487-490.

Gandhi.S, Dhavale.N.D, Jadhav.V.Y, Sabnis.S.S,(2008), Spectrophotometric and RP-HPLC methods for simultaneous determination of escitalopram oxalate and clonazepam in combined tablet dosage form, J.AOAC Int.,91(1),33-38.

Greiner.C, Hiemke.C, Bader.W, Haen.E,(2007), Determination of citalopram and escitalopram together with their active main metabolites desmethyl(es-)citalopram in human serum by column-switching HPLC and spectrophotometric detection, J.Chromatogra.B.,848(2),391-394.

Hemamrutha.S, Rambabu.R, Vidhyadhara.S,(2013), Development and validation of RP-HPLC method for simultaneous estimation of irbesartan and hydrochloro thiazide in in bulk and pharmaceutical dosage forms, Int.J Pharm.,3(2),360-363.

Hermann.M, Christensen.H, Rebsaet.J.L,(2005), Determination of atorvastatin and metabolites in human plasma with solid phase extraction followed by LC-tandem MS, Anal.Bioanal.Chem.382,1242-1249.

Hisao.K, Hajimu.K, Ryozo.I,(1999), HPLC determination of gatifloxacin, a new quinolone, in biological fluids, Japanese J.Chemotherapy,47(2),104-111.

Kabra.P.V, Kimbahune.R.V, Puranik.M.P,(2010), Simultaneous estimation of gatifloxacin,ornidazole and it’s isomers by RP-HPLC, Int.J.Chem.Tech.Res.,2(2), 928-931.

Kahsay.G, Schepdael.A.V, Adams.E,(2012), Development and validation of a liquid chromatographic method for purity control of clopidogrel-acetylsaicylic acid in combined oral dosage forms, J.Pharm.Biomed.Anal.,61,271-276.

Kakde.R.B, Satone.D.D,(2009), Spectrophotometric method for simultaneous estimation of escitalopram oxalate and clonazepam in tablet dosage form, Indian J.Pharm.Sci.71(6),702-705.

Krishna.K.B, Mahendra.K, Sundar.S.B,(2011), A validated RP-HPLC method for the simultaneous estimation of irbesartan and hydrochlorothiazide in pharmaceutical dosage forms, Der Pharma Chemica,3(1),490-496.

Kumar.P.M, Sreeramulu.J,(2011), Development and validation of a stability indicating RP-HPLC method for assay of irbesartan in pure and pharmaceutical dosage form, Int J PharmaSci Rev Res., 6(1),94-99.

Kurien.J, Jayasehar.P,(2013), Stability indicating HPLC determination of clopidogrel bisulfate in pharmaceutical dosage forms, Int. J Compre.Pharma, 1(06), 1-5.

Mitakos.A, Panderi.I,(2002), A validated LC method for the determination of clopidogrel in pharmaceutical preparations, J Pharm.Biomed.Anal.,28(3-4),431-438.

Nagarjuna.A, Rghavacharyulu.K.S.V, Bindu.H.K, Mukkanti.K, Surynarayana.M, (2006), An isocratic chiral sensitive HPLC method developed for the separation of escitalopram oxalate drug substance, Indian drugs,43,746.

Overholser.B.R, Kays.M.B, Sowinski.K.M,(2003), Determination of gatifloxacin in human serum and urine by HPLC with ultra violet detection, J.Chromatogr.B, 798,167-173.

Paramane.S,Kothapalli.L, Thomas.A, Deshpande.A,(2007), Simultaneous RP-HPLC estimation of gatifloxacin and ornidazole in tablet dosage forms, Indian J. Pharma.Sci.69(4),525-528.

Peer.C.J, Spencer.S.D, VanDenBerg.D.A, Pacanowski.M.A, Horenstein.R.B, Figg.W.D,(2012), A sensitive and rapid UPLC-MS/MS method for the simultaneous detection of clopidogrel and it’sderivatized active thiol metabolite in human plasma, J Chromatogr B,880(1),132-139.

Prashanthi.R, Raghavi.K, Sindhura.M, Anupama.B, Nalluri.B.N,(2012), Development and validation of a sensitive RP-HPLC-PDA method for assay of irbesartan in pure and pharmaceutical dosage forms, Int J Pharma.Bio.Sci.,3(1), 397-406.

Prathap.B, Dey.A, Rao.S.G.H, Johnson.P, Arthanariswaran.P,(2013), A Review-Importance of RP-HPLC in Analytical Method Development. Int J Novel Trends Pharm Sci.,3(1),13-23.

Raju.B.V, Rao.L.A,(2011), Validated RP-HPLC method for the estimation of irbesartan in bulk and tablet dosage form, Int J Res Pharma Chem.,1(1),25-28.

Ramakrishna.N, Srinivasalu.K, Saritha.V, Bujjibabu.L, Vishwottam.K, Koteshwara.M, Bindu.H.V,(2010), HPLC method for the separation of enantiomericgatifloxacin, J.Chromatogr.Sci.,48,100-103.

Raman.B, Sharma.B.A, Ghugare.P.D, Karumuse.P.P, Kumar.A,(2009), Semi-preparative isolation and structural elucidation of an impurity in citalopram by LC/MS/MS, J.Pharma.Biomed.Anal.,50(3),377-383.

Raman.B, Sharma.B.A, Ghusare.P.D, Nandavadekar.S, Singh.D, Karmuse.P.K, Kumar.A,(2010), Structural elucidation of process-related impurities in escitalopram by LC/ESI-MS and NMR, J.Pharm.Biomed.Anal.,53(4),895-901.

Rane.V.P, Patil.K.R, Sangshetti.J.N, Yeole.R.D, Shinde.B,(2010), Stability indicating LC method for simultaneous determination of irbesartan and hydrochlorothiazide in pharmaceutical preparations, J Chromatogra. Sci.48,595-600.

Rao.R.N, Raju.A.N, Narsimha.R,(2012), Isolation and characterization of degradation products of citalopram and process-related impurities using RP-HPLC, J.Sep.Sci.31(10),1729-1738.

Reddy.R, Kumar.GVS, Puranik.S.B, Giriprasad.P, Sridhar.K.A,(2012), Development and validation of stability indicating RP-HPLC- method for simultaneous estimation of irbesartan and hydrochlorothiazide in bulk drug and tablet dosage form, Int.J Pharma.Chem.Bio.Sci.2(4),696-703.

Salgado.H.R, Lopes.C.C.G.O,(2006), Determination of gatifloxacin in bulk and tablet preparations by HPLC,89,642-645.

Samanta.T, Dey.S, Bhusansamal.H, Bharatkumar.D, Mohanty.D.L, Kausikbhar,(2011), RP-HPLC method for the estimation of escitalopram in bulk and in dosage forms, Int.J.Chem.Res.,2(2),11-15.

Santoro.M, Kassab.N.M, Singh.A.K, Kedor-Hackam.E,(2006), Quantitative determination of gatifloxacin,levofloxacin,lomefloxacin and pefloxacinfluoroquinolonic antibiotics in pharmaceutical preparations by HPLC, J.Pharm. Biomed.Anal.,40,179-184.

Sharma.D, Mittal.R, Gupta.A, Singh.K, Anroop.N,(2010), Quantitative Bioanalysis by LC-MS/MS: A Review. J Pharma Biomed Sci.7(01),1-9.

Silvestro.L, Gheorghe.M.C, Tarcomnicu.I, Savu.S, Savu.S.R, Lordachescu.A, Dulea.A,(2010), Development and validation of an HPLC-MS/MS method to determine clopidogrel in human plasma. Use of incurred samples to test back-conversion. J chromatography B,878(30),3134-3142.

Singh.S.S, Sha.H, Guptha.S, Jain.N, Sharma.K, Thakkar.P,(2004), Liquid chromatography electrospray ionization mass spectrometry method for the determination of escitalopram in human plasma and it’s application in bio-equivalence study, J.Chromatogr.B.,811,209-215.

Sridhar.S, Divya.S, Madhuri.R, Sudhakar.M,(2013), UPLC- A dynamic and expeditious approach to liquid chromatography. Int J Pharm Chem Bio Sci, 3(4), 1139-1152.

Srinivas.G, Kumar.K.N, Chandrasekhar.R, Ashok.E.D, Rakeshwar.B,(2011), Identification,synthesis and spectral characterization of impurities inprocess development of escitalopram,GERF Bulletin of Biosci.2(1),1-5.

Srinivas.P, Rao.S.T, Ramakrishnan.E, Gowda.D.V,(2013), Development and validation of analytical method for the estimation of omeprazole,domperidone and their substance, Int J Pharm Bio Sci,4(2),906-911.

Srivastava.P.K, Basniwal.P.K, Jain.D, Srivastava.S.K,(2008), Concurrent estimation of clopidogrel bisulfate and aspirin in tablets by validated RP-HPLC method, Ind.J Pharm.Sci.,70(5),667-669.

Sundar.S.B,(2011), Development and validation of liquid chromatographic method for estimation of escitalopram oxalate in tablet dosage forms, Int.J.Pharm. Bio.Sci., 2(1),140-146.

Tadic.S, Nikolic.K, Agbada.D,(2012), Development and optimization of an HPLC analysis of citalopram and it's four nonchiral impurities using experimental design methodology, J.AOAC Int.,95(3),733-743.

Vetrichelvan.T, Arul.K, Sumithra.M, Umadevi.B,(2010), Colorimetric method for the estimation of escitalopram oxalate in tablet dosage form, Indian J.Pharm.Sci.,72(2),269-271.

Vukkum.P, Babu.J.M, Muralikrishna.R,(2013), Stress degradation behavior of atorvastatin calcium and development of a suitable stability-indicating LC method for the determination of atorvastatin, its related impurities and its degradation products, Sci.Pharm.,81(1),93-114.